arXiv:cond-mat/0003366v1 [cond-mat.stat-mech] 22 Mar 2000

Comment on "Anomalous Spreading of Power-Law Quantum Wave Packets"

Anomalous spreading of free wave packets with power-law tails was reported in a recent Letter by Lillo and Mantegna [1]. The authors consider positive even wave functions $\psi(x)$ decreasing with x as $\psi(x) \sim |x|^{-\alpha}$ ($\alpha > 1/2$), and analyze the asymptotic time evolution through the stationary phase approximation. They conclude that, for $\alpha > 1$ the well known upper bound $t^{-1/2}$ for the decay of Gaussian and finite extension packets still stands, while for $1/2 < \alpha < 1$ the decay follows the $t^{-\alpha/2}$ law. However, they do not give a correct answer for the case $\alpha = 1$. As an illustrative example they work out the special family of wave functions

$$\psi(x) = N\,(x^2+\gamma^2)^{-\alpha/2}, \tag{1}$$

where N is a normalization constant and γ a real parameter. This family naturally emerges, in fact, in anomalous diffusion [2].

We point out that their main result, i.e., the anomalous decay of wave functions with $|x|^{-\alpha}$ tails was derived before in Ref. [3] where a wave function similar to (1) was used as prototype.

A result for general functions decaying as $|x|^{-1}$ is not provided in [1]. However, the authors state that, in the special case described by Eq. (1) and $\alpha = 1$, the maximum of the packet decreases as $1/\sqrt{t}$, *an assertion which is not correct.*

We want to show that there is an extremely direct way to obtain the *exact* decay of packets given by Eq. (1). In particular, it is shown that $\alpha = 1$ leads to a $\ln(t)/\sqrt{t}$ decay, as already pointed out in [4]. As it is well known, the time evolution of a free wave packet in one dimension can be described by [5]

$$\psi(x,t) = \frac{1}{\sqrt{2\pi i\,\hbar t/M}} \int_{-\infty}^{\infty} \mathrm{e}^{\frac{iM\left(x-x'\right)^2}{2\hbar t}}\, \psi(x',0)\,\mathrm{d}x', \tag{2}$$

where M is the mass of the particle. Following Ref. [1], we concentrate our attention in the behavior of the maximum of the wave packet (at $x = 0$). Then, we realize that the integral in Eq. (2) when $\psi(x,0)$ is given by Eq. (1) can be evaluated directly by using the following integral representation of the Kummer function, $U(\nu,\mu,z)$ [6], with $\nu = \frac{1}{2}$ and $\mu = \frac{3-\alpha}{2}$,

$$U(\tfrac{1}{2},\tfrac{3-\alpha}{2},z) = \frac{1}{\sqrt{\pi}} \int_0^{\infty} \mathrm{e}^{-zs} s^{-1/2} (1+s)^{-\alpha/2} \mathrm{d}s. \tag{3}$$

Hence, $\psi(0,t)$ can be obtained straightforwardly from Eq. (3) after the change of variables $s = (x'/\gamma)^2$ together with the identification $z = \gamma^2 M/[2i\hbar t]$. In this way we get

$$\psi(0,t) = \frac{N\gamma^{1-\alpha}}{\sqrt{2i\hbar t/M}}\, U(\tfrac{1}{2},\tfrac{3-\alpha}{2},\gamma^2 M/[2i\hbar t]). \tag{4}$$

In order to obtain the asymptotic time behavior of $\psi(0,t)$, we employ the following limiting forms of the Kummer function for small $|z|$ [7]

$$U(\tfrac{1}{2},\tfrac{3-\alpha}{2},z) \simeq \begin{cases} \frac{1}{\sqrt{\pi}}\Gamma(\frac{1-\alpha}{2})\, z^{(\alpha-1)/2} & (1/2<\alpha<1), \\ -\frac{1}{\sqrt{\pi}}\, \ln(z) & (\alpha=1), \\ \Gamma(\frac{\alpha-1}{2})/\Gamma(\frac{\alpha}{2}) & (1<\alpha). \end{cases} \tag{5}$$

Finally, replacing these asymptotic expressions in Eq. (4) we get for large t

$$|\psi(0,t)| \sim \begin{cases} t^{-\alpha/2} & (1/2<\alpha<1), \\ \ln(t)\, t^{-1/2} & (\alpha=1), \\ t^{-1/2} & (1<\alpha). \end{cases} \tag{6}$$

So, the $\ln(t)/\sqrt{t}$ law is expected to be the general one governing the asymptotic evolution of packets decaying as $|x|^{-1}$ since tails and not other details of the packet shape seem to rule the asymptotic behavior.

We acknowledge partial financial support from Brazilian research agencies FAPERJ and CNPq.

R. S. Mendes[1,2] and C. Anteneodo[3]

[1]Departamento de Física, Universidade Estadual de Maringá, Av. Colombo 5790, CEP 87290-020, Maringá-PR, Brazil, e-mail: rsmendes@dfi.uem.br

[2]Department of Physics, University of North Texas, Denton, TX 76203, USA

[3]Instituto de Biofísica, Universidade Federal do Rio de Janeiro, C. Universitária, CCS, CEP 91949-900, Rio de Janeiro, Brazil, e-mail: celia@cat.cbpf.br

PACS numbers: 03.65.-w, 05.60.+w

---

[1] F. Lillo and R. N. Mantegna, Phys. Rev. Lett. **84** 1061 (2000).

[2] C. Tsallis et al, Phys. Rev. Lett. **75**, 3589 (1995) [Erratum: **77**, 5442 (1996)]; A.R. Plastino and A. Plastino, Physica A **222**, 347 (1995); C. Tsallis and D.J. Bukman, Phys. Rev. E **54**, R2197 (1996).

[3] K. Unnikrishnan, Am. J. Phys. **66**, 632-633 (1998).

[4] K. Unnikrishnan, Am. J. Phys. **65**, 526-527 (1997).

[5] K. Gottfried, *Quantum mechanics* (W. A. Benjamin, Inc., Amsterdam, 1966), p. 22.

[6] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series 55 (Washington, 1965), p. 505.

[7] *Ibid.*, p. 508.